

Alameda County Health Care Services Agency Public Health Department

www.acphd.org

Colleen Chawla, Director Kimi Watkins-Tartt, Director Nicholas Moss, MD, Health Officer

Public Health Department: Main Line (510) 267-8000

COVID-19 Information: (510) 268-2101

MPX Isolation Guidance

Last Updated: September 21, 2022

PURPOSE OF THIS DOCUMENT

The purpose of this document is to provide MPX isolation guidance. The following recommendations apply to persons with laboratory confirmed MPX, as well as persons with MPX symptoms while awaiting testing or lab results (i.e., suspected MPX).

If you suspect you may have MPX, isolate and contact your healthcare provider immediately. If you do not have a regular medical provider, contact us at 510-268-2101 or Monkeypox@acgov.org for support.

MPX OVERVIEW

MPX—also known as Monkeypox—is a rare disease caused by infection with the MPX virus. While this virus is related to smallpox, MPX illness is milder than smallpox but can still be serious.

HOW MPX SPREADS

Any person—of any age, gender, or sexual orientation—who has direct physical contact with an infected person can get MPX. **The MPX virus can spread by:**

- Direct skin-to-skin contact with the rash, sores, scabs, or body fluids of a person with infection
- Sexual and other intimate contact (e.g., hugging, cuddling, and kissing) with a person with infection
- Sharing unwashed bedding, towels, clothing, and utensils used by a person with infection
- Prolonged and close face-to-face contact with someone with infection
- Passing from a pregnant mother to the fetus

SYMPTOMS OF MPX

The most common symptom of MPX is a **new, unexplained rash** that can look like pimples or blisters on the face, inside the mouth, and on other parts of the body, like the hands, feet, chest, genitals, and around the anus.

Early or additional symptoms of MPX may include **flu-like symptoms**, such as:

- Fever and/or Chills
- Headache
- Muscle and/or Body Aches
- Swollen Lymph Nodes
- Tiredness and/or Weakness

Note: If present, flu-like symptoms usually appear about 5 days before a rash.



Alameda County Health Care Services Agency Public Health Department

www.acphd.org

Colleen Chawla, Director Kimi Watkins-Tartt, Director Nicholas Moss, MD, Health Officer

Public Health Department: Main Line (510) 267-8000

COVID-19 Information: (510) 268-2101

ISOLATION INSTRUCTIONS

In accordance with the California Public Health Department (CDPH), persons with confirmed or suspected MPX should isolate (stay at home in a private room) unless it is necessary to see a healthcare provider.

What is Isolation?
Any person with suspected or confirmed MPX should isolate, immediately. It is best to isolate in a single room
with a dedicated bathroom
☐ Avoid close physical contact with other people and animals
☐ Keep rash and open sores completely covered with clothing, bandages, and/or gloves
☐ Wear a well-fitting mask when around other people
☐ Do not share beds, clothing, sheets, blankets, towels, washcloths, or eating utensils
☐ Wash hands or use alcohol-based hand sanitizer frequently
 Clean and disinfect all surfaces that you touch in shared spaces (especially the bathroom).
Note : It is okay to share spaces with others who are infected with MPX.
For more information on MPX prevention, see the following state and federal guidance:
 MPX Home Isolation Guidance for the General Public (CDPH)
 <u>Isolation and Infection Control at Home (CDC)</u>
 <u>Disinfection of the Home and Non-Healthcare Settings (CDC)</u>
How long should I isolate?
Persons with MPX can transmit infection from the start of MPX symptoms until all sores have fully healed and a
fresh layer of skin has formed. This process can take 2-4 weeks. This is called the "infectious period"
If you tested positive for MPX, you should isolate at home until meeting <u>all</u> the following criteria:
☐ No fever (temperature of ≥100.4°F) for at least 48 hours without use of fever-reducing medication; and
☐ No respiratory symptoms (e.g., sore throat, nasal congestion, and cough) for at least 48 hours; and
Two respiratory symptoms (e.g., sore timout, masar congestion, and coagil, for at least 40 hours, and
☐ No new lesions have appeared for at least 48 hours; and
 □ No new lesions have appeared for at least 48 hours; and □ Any rash and all sores have scabbed and healed with a fresh layer of skin, including sores in the mouth.
 □ No new lesions have appeared for at least 48 hours; and □ Any rash and all sores have scabbed and healed with a fresh layer of skin, including sores in the mouth. You can resume some activities outside the home that do not involve higher risk settings* or physical contact
 □ No new lesions have appeared for at least 48 hours; and □ Any rash and all sores have scabbed and healed with a fresh layer of skin, including sores in the mouth. You can resume some activities outside the home that do not involve higher risk settings* or physical contact with other people if all the following criteria are met:
 □ No new lesions have appeared for at least 48 hours; and □ Any rash and all sores have scabbed and healed with a fresh layer of skin, including sores in the mouth. You can resume some activities outside the home that do not involve higher risk settings* or physical contact

RETURNING TO WORK



Alameda County Health Care Services Agency Public Health Department

www.acphd.org

Colleen Chawla, Director Kimi Watkins-Tartt, Director Nicholas Moss, MD, Health Officer

Public Health Department: Main Line (510) 267-8000

COVID-19 Information: (510) 268-2101

If you work in a **higher risk setting** you should wait to return to work until any rash and all sores have scabbed and fully healed with a fresh layer of skin. Please refer to <u>CDPH MPX Home Isolation Guidance for the General Public, Section C</u> for detailed instructions on when it is safe for you to return to work.

*Higher Risk Settings include:

- Homeless shelters, migrant shelters, emergency shelters, and residential drug treatment facilities
- Healthcare settings
- State and local correctional facilities and detention centers.
- Long-term care, adult and senior care facilities, and in-home services involving physical care
- Childcare and preschool settings that provide care for children from infancy through pre-school, assuming that there will be close physical contact for diapering, toileting, feeding, hygiene and general interaction
- K-12 schools and other settings (before/after school programs) that provide care for school-aged children younger than age 8 or older children whose care requires close physical contact, if job duties require direct physical contact with such children

NOTIFY YOUR CONTACTS

Make sure to tell anyone you had close physical contact with since your symptoms started that you have MPX infection if it is safe for you to do so. Include anyone with whom you had sex, cuddled, kissed, or shared a bed/room. People who have been exposed may be offered vaccination as soon as possible to prevent MPX illness or reduce symptoms. Exposed contacts can go to <u>Alameda County Public Health Department MPX Vaccine</u> webpage to find out where they can get vaccinated.

ANSWER CALLS from PUBLIC HEALTH

Public health nurses and investigators are working to limit the spread of MPX. They will talk to you about your symptoms, ask about your contacts, and offer you resources to support your health and well-being.

ADDITIONAL IMPORTANT LINKS

Alameda County MPX Website

Centers for Disease Control and Prevention: If You Are Sick

California Department of Public Health: MPX homepage

California Department of Public Health: Home Isolation Guidance for the General Public